



TITLE:

Log-canonical models of singular pairs and its applications

AUTHOR(S):

Odaka, Yuji; Xu, Chenyang

CITATION:

Odaka, Yuji ...[et al]. Log-canonical models of singular pairs and its applications. Mathematical Research Letters 2012, 19(2): 325-334

ISSUE DATE:

2012

URL:

<http://hdl.handle.net/2433/160402>

RIGHT:

© International Press of Boston, Inc.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

LOG-CANONICAL MODELS OF SINGULAR PAIRS AND ITS APPLICATIONS

YUJI ODAKA AND CHENYANG XU

ABSTRACT. We prove the existence of the log-canonical model over a log pair (X, Δ) . As an application, together with Kollár’s gluing theory, we remove the assumption in the first named author’s work [Odaka11], which shows that K -semistable polarized varieties can only have semi-log-canonical singularities.

CONTENTS

1. Introduction
 2. Log-canonical models
 3. Semi-log-canonical models
- Acknowledgment
References

1. Introduction

Throughout this paper, the ground field is assumed to be an algebraically closed field of characteristic 0. It is well known that a normal surface singularity has the *minimal resolution*, while for a singular variety of higher dimension, usually it does not have any such “canonically determined” smooth modification. But if we allow the partial resolution having mild singularities, a type of singularities coming from the minimal model program (MMP) which is natural for many questions, then it is possible. More precisely, for an arbitrary normal variety X , we can consider a unique (“canonically determined”) partial resolution $Y \rightarrow X$ with only canonical singularities and satisfies the property that K_Y is relative ample over X . The existence of such model Y , i.e., *the canonical model over X^1* , is implied by [BCHM10, Main theorem (1.2)]. In the case of surfaces, Y is obtained by contracting all exceptional curves with self-intersection (-2) from the minimal resolution.

Similarly, for a normal pair (X, Δ) i.e., attached with a boundary \mathbb{Q} -divisor, we can define a “canonically determined” partial resolution $(Y, \Delta_Y) \rightarrow (X, \Delta)$ associated to it, which is called its *log-canonical model* (see (2.1)). It coincides with the relative log-canonical model of a log resolution with a reduced boundary, in the sense of the usual relative log MMP, as we will show in Lemma 2.1.

In this note, we study the question of the existence of log-canonical model of a normal pair (X, Δ) . It is well known that the full log MMP (including the abundance

Received by the editors October 3, 2011.

¹Here, the adjective “canonical” comes from the sense of singularities.

conjecture) gives an affirmative answer to the question. As the full log MMP is still not established, our main observation in this note is that if we assume $K_X + \Delta$ is \mathbb{Q} -Cartier, then the existence of log-canonical model follows from the established results on MMP, especially the recent ones in [Birkar11] and [HX11].

Theorem 1.1. *Let (X, Δ) be a normal pair, i.e., X is a normal variety and $\Delta = \sum a_i \Delta_i$ is a \mathbb{Q} -divisor with distinct prime divisors Δ_i and rational numbers a_i . Assume $0 \leq a_i \leq 1$ and $K_X + \Delta$ is \mathbb{Q} -Cartier. Then there exists a log-canonical model (Y, Δ_Y) over (X, Δ) (see (2.1) for the definition).*

As a consequence, we give a proof of the inversion of adjunction for log canonicity, which is a slight simplification of Hacon's argument in [Hacon11] (also see [Kollár12, 4.11.2]). We note that the inversion of adjunction for log canonicity was first proved by Kawakita (see [Kawakita07]) without using the minimal model program.

Corollary 1.1 (Inversion of Adjunction). *Let $(X, D + \Delta)$ be a normal pair and D a reduced divisor. Assume $K_X + D + \Delta$ is \mathbb{Q} -Cartier. Let $n: D^n \rightarrow D$ be the normalization, and write $n^*(K_X + D + \Delta)|_{D^n} = K_{D^n} + \Delta_{D^n}$.*

Then $(X, D + \Delta)$ is log-canonical along D if and only if (D^n, Δ_{D^n}) is log-canonical.

We can also extend our results into non-normal setting. In fact, Kollár recently has developed a rather complete theory of semi-log-canonical pairs by studying their normalizations. Thanks to his fundamental theory (see [Kollár12]), including his recent result [Kollár11], we have the following as a consequence, which generalizes Theorem 1.1.

Corollary 1.2. *Let (X, Δ) be a demi-normal pair where $\Delta = \sum a_i \Delta_i$ is a \mathbb{Q} -divisor, none of prime divisor Δ_i are in the singular locus $\text{Sing}(X)$. Assume $0 \leq a_i \leq 1$ and $K_X + \Delta$ is \mathbb{Q} -Cartier. Then the semi-log-canonical model (Y, Δ_Y) over (X, Δ) exists.*

Recall that *demi-normality* of X means that it is normal crossing in codimension 1 and satisfies Serre's S_2 condition [Kollár12, 5.1]. For the precise definition of *semi-log-canonical model*, see Definition 3.2.

One of our main applications for this note is the following. In [Odaka11], the first named author proved K -semi-stability implies semi-log canonicity, assuming the existence of semi-log-canonical models. Since (1.2) verifies this assumption, the following theorem now becomes unconditional.

Theorem 1.2 ([Odaka11]). *Let X be an equidimensional reduced projective variety, satisfies S_2 condition and whose codimension 1 points are Gorenstein. Thus we can define the Weil divisor class K_X which we assume to be \mathbb{Q} -Cartier.*

Then, if (X, L) is K -semistable, X has only semi-log-canonical singularities.

Roughly speaking, assuming the non-semi-log-canonicity of X , Odaka [Odaka11] proved that we can construct “destabilizing test configuration” by using the semi-log-canonical model of X . We refer to [Odaka11] for more details.

2. Log-canonical models

Definition 2.1. Let (X, Δ) be a normal pair, i.e., X is a normal variety and $\Delta = \sum a_i \Delta_i$ is a \mathbb{Q} -divisor with distinct prime divisors Δ_i and rational numbers a_i . Assume

$0 \leq a_i \leq 1$. We call that a birational projective morphism $f : Y \rightarrow (X, \Delta)$ give a *log-canonical model* over (X, Δ) if with the divisor $\Delta_Y = f_*^{-1}(\Delta_X) + E_f^{\text{lc}}$ on Y , where E_f^{lc} denotes the sum of f -exceptional prime divisors with coefficients 1, the pair (Y, Δ_Y) satisfies

- (1) (Y, Δ_Y) is log-canonical,
- (2) $K_Y + \Delta_Y$ is ample over X .

From the negativity lemma (see [KM98, 3.38]), we know that $f : Y \rightarrow X$ is isomorphic over the maximal open locus X^{lc} on which (X, Δ) is log-canonical (see the proof of (2.2)). For more background of log-canonical models over a pair (X, Δ) , see [Koll retal92, Section 2].

First, we discuss the uniqueness of the log-canonical model.

Lemma 2.1. *Let $\tilde{f} : \tilde{Y} \rightarrow X$ be a log resolution of (X, Δ) . Assume that $(\tilde{Y}, \Delta_{\tilde{Y}} := \tilde{f}_*^{-1}\Delta + \sum E_i)$ has a relative log-canonical model (Y, Δ_Y) over X , where E_i run over all \tilde{f} -exceptional prime divisors. Then $Y \rightarrow (X, \Delta)$ is a log-canonical model over (X, Δ) .*

Proof. By the definition of the relative log-canonical model, (Y, Δ_Y) obviously satisfies conditions (1) and (2). \square

Proposition 2.1. *If log-canonical model Y exists, then it is unique.*

Proof. Let $g : \tilde{Y} \rightarrow Y$ be a log resolution of $(Y, f_*^{-1}(\Delta) + \text{Ex}(f))$. And we write

$$g^*(K_Y + \Delta_Y) + E \sim_{\mathbb{Q}} K_{\tilde{Y}} + F,$$

such that $E, F \geq 0$, have no common components. It is easy to see that $g_*(F) = \Delta$. Since (Y, Δ_Y) is log-canonical

$$\tilde{f}_*^{-1}\Delta + \sum E_i \geq F,$$

where $\tilde{f} = f \circ g$ and E_i run over all \tilde{f} -exceptional prime divisors. The difference is g -exceptional. We conclude that

$$\text{Proj} \bigoplus_{m \in \mathbb{Z}_{\geq 0}} \tilde{f}_* \mathcal{O}_{\tilde{Y}}(m(K_{\tilde{Y}} + \Delta_{\tilde{Y}})) \cong \text{Proj} \bigoplus_{m \in \mathbb{Z}_{\geq 0}} f_* \mathcal{O}_Y(m(K_Y + \Delta_Y)) \cong Y,$$

as $K_Y + \Delta_Y$ is ample over X . So it suffices to show that the different log resolutions as in (2.1) will yield the same log-canonical model Y .

We assume that there are two difference choices $g_i : (\tilde{Y}_i, \Delta_{\tilde{Y}_i}) \rightarrow (X, \Delta)$ ($i = 1, 2$) with a morphism $\mu : \tilde{Y}_1 \rightarrow \tilde{Y}_2$. Since $\mu^*(K_{\tilde{Y}_2} + \Delta_{\tilde{Y}_2}) + E' = K_{\tilde{Y}_1} + \Delta_{\tilde{Y}_1}$ for some effective exceptional divisor E' . The uniqueness immediately follows from the fact that $(\tilde{Y}_i, \Delta_{\tilde{Y}_i})$ have the same relative log-canonical ring (sheaf)

$$\bigoplus_{m \in \mathbb{Z}_{\geq 0}} (g_i)_* \mathcal{O}_{\tilde{Y}_i}(m(K_{\tilde{Y}_i} + \Delta_{\tilde{Y}_i}))$$

over X . \square

Lemma 2.2. *Let (X, Δ) be a pair as in (2.1). We assume that $K_X + \Delta$ is \mathbb{Q} -Cartier. Let $f : Y \rightarrow (X, \Delta)$ be the log-canonical model. Write*

$$f^*(K_X + \Delta) \sim_{\mathbb{Q}} K_Y + B,$$

and $B = \sum b_i B_i$ as the sum of distinct prime divisors such that $f_(B) = \Delta$, we let $B^{>1}$ be the nonzero divisor $\sum_{b_i > 1} b_i B_i$ and $B^{\leq 1}$ be the divisor $\sum_{b_i \leq 1} b_i B_i$, then $\text{Supp}(B^{>1}) = \text{Ex}(f)$. In particular, $\text{Ex}(f) \subset Y$ is of pure codimension 1.*

Proof. It is obvious that $\text{Supp}(B^{>1}) \subset \text{Ex}(f)$.

If we write $B = f_*^{-1}(\Delta) + E_B$, then E_B is supported on the exceptional locus and the divisor $E_f^{\text{lc}} - E_B$ is an exceptional divisor which is relatively ample. It follows from the negativity lemma (see [KM98, 3.38]) that $E_f^{\text{lc}} - E_B \leq 0$. Therefore, we have the equality

$$f_*^{-1}(\Delta) + E_f^{\text{lc}} = B^{\leq 1} + \text{Supp}(B^{>1}).$$

From the definition of the log-canonical model (2.1), we know that

$$(2.1) \quad K_Y + \text{Supp}(B^{>1}) + B^{\leq 1} \sim_{\mathbb{Q}, X} \text{Supp}(B^{>1}) - B^{>1} = \sum_{b_i > 1} (1 - b_i) B_i$$

is relatively ample over X . Thus for any curve C which is contracted by f , we have

$$C \cdot \left(\sum_{b_i > 1} (b_i - 1) B_i \right) < 0,$$

which implies that $C \subset \text{Supp}(B^{>1})$. This shows $\text{Ex}(f) \subset \text{Supp}(B^{>1})$ which completes the proof. \square

Proof of (1.1). We take a (\mathbb{Q} -factorial) dlt modification $g : Z \rightarrow X$ of (X, Δ) (see [KK10, Section 3] or [Fujino10, 4.1]) such that

- (i) if we write $g^*(K_X + \Delta) \sim_{\mathbb{Q}} K_Z + g_*^{-1}(\Delta) + \sum b_i E_i$, then $b_i \geq 1$;
- (ii) $(Z, \Delta_Z = g_*^{-1}(\Delta) + \sum E_i)$ is dlt.

We remark Z can be achieved by running a sequence of $(K_{\tilde{Y}} + \Delta_{\tilde{Y}})$ -MMP over X for a log resolution of $\tilde{f} : \tilde{Y} \rightarrow (X, \Delta)$ (see [Fujino10, 4.1]) where $\Delta_{\tilde{Y}}$ is defined as in (2.1). Furthermore, we require that $\tilde{f}^{-1}(X \setminus X^{\text{lc}})$ is a divisor. We want to show $(Z, \Delta_Z = g_*^{-1}(\Delta) + \sum E_i)$ has a good minimal model over X . Since then we can take Y to be the relative log-canonical model, which is easy to see it is the log-canonical model of (X, Δ) .

Lemma 2.3. *Let V be a log-canonical center of $(Z, g_*^{-1}(\Delta) + \sum E_i)$, such that $g(V) \subset X \setminus X^{\text{lc}}$, then $V \subset E_i$ for some i .*

Proof. As Z is obtained by running a sequence of MMP for a log smooth resolution $\tilde{f} : \tilde{Y} \rightarrow (X, \Delta)$, then V is an lc center of $(Z, g_*^{-1}(\Delta) + \sum E_i)$ if and only if $\tilde{Y} \dashrightarrow Z$ is isomorphic over the generic point V and the preimage W of V in \tilde{Y} is a component of $\cap F_i$, where F_i 's are prime divisors contained in $\lfloor \Delta_{\tilde{Y}} \rfloor$. As by our assumption $\tilde{f}^{-1}(X \setminus X^{\text{lc}})$ is a union of divisors, if $\tilde{f}(W) \subset X \setminus X^{\text{lc}}$, then W is contained in one of the \tilde{f} -exceptional divisors \tilde{E}_i whose image is in $X \setminus X^{\text{lc}}$. Therefore, V is contained the

birational transform of \tilde{E}_i on Z as $\tilde{Y} \dashrightarrow Z$ is an isomorphism on the generic point of \tilde{E}_i . \square

Now consider all exceptional divisors E of g with the centers contained in $X \setminus X^{\text{lc}}$. Fixing a general relatively ample effective divisor H on Z over X , we run $(K_Z + \Delta_Z)$ -MMP with scaling of H over X (see [BCHM10, Subsection 3.10]). As we treat dlt pairs, which are not klt, we explain what follows from [BCHM10] in the following lemma for the readers' convenience.

Lemma 2.4. *We can run the MMP with scaling of H for (Z, Δ_Z) over X to get a sequence of numbers $0 \leq \cdots \leq s_2 \leq s_1 \leq s_0$ and a sequence of birational models*

$$Z = Z_0 \dashrightarrow Z_1 \dashrightarrow Z_2 \dashrightarrow \cdots,$$

such that the following holds. Here, Δ_j and H_j are push-forwards of Δ and H on each Z_j .

- (i) $K_{Z_j} + \Delta_j + tH_j$ is semi-ample over X for any $s_j \geq t \geq s_{j+1}$.
- (ii) This sequence $\{s_i\}$ (is either finite with $\exists s_N = 0$ or) satisfies the property that $\lim_j s_j = 0$.

Proof. For each Z_j , we set

$$s_{j+1} := \inf\{t > 0 \mid K_{Z_j} + \Delta_j + tH_j \text{ is relatively nef over } X\},$$

and consider the extremal contraction of an extremal ray R_j with $(K_{Z_j} + \Delta_j + s_{j+1}H_j) \cdot R_j = 0$. In each step the existence of flip holds since as $K_{Z_j} + \Delta_j$ is dlt and Z_j is \mathbb{Q} -factorial, $(K_{Z_j} + \Delta_j)$ -flip is the same as a step of $(K_{Z_j} + (1 - \delta)\Delta_j)$ -MMP for $0 < \delta \ll 1$ and $(Z_j, (1 - \delta)\Delta_j)$ is klt (see [BCHM10, Corollary 1.4.1]).

From our construction, we know that giving a sequence of j steps

$$Z = Z_0 \dashrightarrow Z_1 \dashrightarrow Z_2 \dashrightarrow Z_j$$

of $(K_Z + \Delta)$ -MMP with scaling of H as above is the same as giving a sequence of steps of $(K_Z + \Delta + tH)$ -MMP with scaling of H for any $0 \leq t < s_j$.

For arbitrary $t > 0$, there exists an effective divisor $\Theta_t \sim_{\mathbb{Q}} \Delta_Z + tH$, such that (Z, Θ_t) is klt (with Θ_t is relatively big, which is trivial in this case since Z is birational over X). It follows from [BCHM10, Corollary 1.4.2, see also Theorem 1.2] that any sequence of $(K_Z + \Theta_t)$ -MMP with scaling of H over X will terminate after finite steps with a relative good minimal model Z_j , i.e., $K_{Z_j} + \rho_{j*}(\Theta_t)$ is semi-ample over X where $\rho_j : Z \dashrightarrow Z_j$ is the birational contraction. (Recall that *good minimal model* means a minimal model which satisfies the abundance conjecture.) Thus, (i) is proved.

Moreover, from the arguments above, there are only finitely many s_j such that $s_j > t$. Since we can choose t to be an arbitrarily small positive number, we also have the conclusion (ii). \square

The *diminished stable base locus*² of (Z, Δ_Z) over Z is defined by

$$\mathbf{B}_-(K_Z + \Delta_Z/X) = \bigcup_{\epsilon > 0} \mathbf{B}(K_Z + \Delta_Z + \epsilon H/X),$$

²Also called *restricted stable base locus*.

where $\mathbf{B}(\cdot)$ denotes the usual stable base locus. If there is a divisor $E \subset \mathbf{B}_-(K_Z + \Delta_Z/X)$, then $E \subset \mathbf{B}(K_Z + \Delta_Z + tH/X)$ for some $t > 0$, therefore there exists an j , such that $s_j \geq t \geq s_{j+1}$. Since

$$K_{Z_j} + \Delta_j + tH_j \sim_{\mathbb{Q}_j} K_{Z_j} + \rho_{j*} \Theta_t$$

is semi-ample over X we know that ρ_j contracts E .

Lemma 2.5. *There exists Z_j such that if we denote by $Z' = Z_j$, $\rho' = \rho_j$, the morphism $g' : Z' \rightarrow X$ and write*

$$g'^*(K_X + \Delta) = K_{Z'} + g'^{-1}(\Delta) + \sum b_i E'_i,$$

then $b_i > 1$ for all E'_i which centers in $X \setminus X^{\text{lc}}$.

Proof. From the above discussion, we can assume that there is $Z_j = Z'$ such that $\mathbf{B}_-(K_{Z'} + g'^{-1}(\Delta) + \sum E'_i)$ has codimension at least 2. By (2.1), we have

$$K_{Z'} + \rho'_*(\Delta_Z) = K_{Z'} + g'^{-1}(\Delta) + \sum E'_i \sim_{\mathbb{Q}, X} \sum (1 - b_i) E'_i,$$

if the statement is not true, it follows from the Kollár–Shokurov’s connectedness theorem (see [Kolláretal92, 17.4]) that there is a divisor E'_0 , with $b_0 = 1$ such that $\sum_{b_i > 1} E'_i|_{E'_0}$ is not trivial. Therefore,

$$\left(\sum (1 - b_i) E'_i + \epsilon H' \right) \Big|_{E'_0}$$

is not effective for small $0 < \epsilon \ll 1$, where $H' := \rho_* H$. This implies that $E'_0 \subset \mathbf{B}_-(K_Z + \Delta_Z/X)$, which yields a contradiction. Then we conclude that $b_i > 1$ for all E'_i whose center is in $X \setminus X^{\text{lc}}$. \square

Now consider the dlt pair $(Z', g'^{-1}(\Delta) + \Sigma)$, where $\Sigma = \sum E'_i - \epsilon \sum (b_i - 1) E'_i$ for some positive $\epsilon \ll 1$.

Lemma 2.6. *$(Z', g'^{-1}(\Delta) + \Sigma)$ has a good minimal model Y' over X .*

Proof. Over the open set X^{lc} , we have

$$(K_{Z'} + g'^{-1}(\Delta) + \Sigma)|_{g'^{-1}(X^{\text{lc}})} = g'^*(K_X + \Delta)|_{g'^{-1}(X^{\text{lc}})},$$

whose ring of pluri-log-canonical sections is finitely generated over X^{lc} , because it is isomorphic to the algebra

$$\oplus_{m \geq 0} \mathcal{O}_{X^{\text{lc}}}(m(K_{X^{\text{lc}}} + \Delta|_{X^{\text{lc}}})) .$$

Therefore, the restriction of $(Z', g'^{-1}(\Delta) + \Sigma)$ over X^{lc} has a relative good minimal model over X^{lc} by [HX11, 2.11]. Any lc center of $(Z', g'^{-1}(\Delta) + \sum E'_i)$ which is contained in on of E'_i cannot be an lc center of $(Z', g'^{-1}(\Delta) + \Sigma)$, however, these lc centers are precisely those centers of $(Z', g'^{-1}(\Delta) + \sum E'_i)$ which is mapped into $X \setminus X^{\text{lc}}$ by (2.3). Thus we conclude that if V is an lc center of $(Z', g'^{-1}(\Delta) + \Sigma)$, then its image under g' intersects X^{lc} . Therefore, it follows from [Birkar11, Theorem 1.9] or [HX11, 1.1] that $(Z', g'^{-1}(\Delta) + \Sigma)$ has a good minimal model $f' : Y' \rightarrow X$. \square

Since

$$K_{Z'} + g'^{-1}_*(\Delta) + \sum E'_i = \frac{1}{1+\epsilon}(K_{Z'} + g'^{-1}_*(\Delta) + \Sigma) + \frac{\epsilon}{1+\epsilon}g'^*(K_X + \Delta),$$

we conclude that Y' is also a relative good minimal model for $K_{Z'} + g'^{-1}_*(\Delta) + \sum E'_i$ over X . \square

Proof of (1.1). One direction is easy (see [Koll retal92, 17.2]). To prove the converse, let us assume that $(X, D + \Delta)$ is not log-canonical along D . Let $f : Y \rightarrow (X, D + \Delta)$ be the log-canonical model as in the proof of (1.1). Write

$$f^*(K_X + D + \Delta) = K_Y + D_Y + B,$$

where D_Y is the birational transform of D . Since f is not an isomorphism over D , it follows from (2.2) that

$$D_Y \cap \text{Ex}(f) = D_Y \cap \text{Supp}(B^{>1}) \neq \emptyset.$$

Therefore, if we denote by D_Y^n the normalization of D_Y and write

$$n^*(K_Y + D_Y + B)|_{D_Y^n} = K_{D_Y^n} + B_{D_Y^n},$$

then $(D_Y^n, B_{D_Y^n})$ has coefficient strictly larger than 1 along some components of $D_Y \cap \text{Ex}(f)$ by (2.2), which implies that (D^n, Δ_{D^n}) is not log-canonical. \square

Corollary 2.1. *Notation as above proof. Let $f : Y \rightarrow (X, D + \Delta)$ be the log-canonical model. Let D_Y be the birational transformation of D and $n : D_Y^n \rightarrow D_Y$ its normalization. Then $f_{D^n} : D_Y^n \rightarrow (D^n, \Delta_{D^n})$ is also the log-canonical model.*

Proof. From the proof of (1.1), we know that

$$n^{-1}(\text{Ex}(f)) = \text{Ex}(f_{D^n}),$$

which implies that if we denote $f_{D^n}^{-1}_*(\Delta_{D^n}) + E_{f_{D^n}}^{\text{lc}}$ by $\Delta_{D_Y^n}$, then

$$K_{D_Y^n} + \Delta_{D_Y^n} = n^*((K_Y + \Delta_Y)|_{D_Y}).$$

Then obviously $(D_Y^n, \Delta_{D_Y^n})$ is log-canonical and $K_{D_Y^n} + \Delta_{D_Y^n}$ is ample over D^n . \square

3. Semi-log-canonical models

In this section, we study the existence of semi-log-canonical model of a demi-normal pair (X, Δ) . A pair (X, Δ) is called *demi-normal* if X is S_2 , whose codimension 1 points are regular or ordinary nodes and Δ is an effective \mathbb{Q} -divisor whose support does not contain any codimensional 1 singular points. For such a demi-normal scheme X , let $n : \bar{X} \rightarrow X$ be its normalization, we can define the *conductor ideal*

$$\text{cond}_X := \text{Hom}_X(n_*\mathcal{O}_{\bar{X}}, \mathcal{O}_X) \subset \mathcal{O}_X.$$

and the *conductor scheme* $D := \text{Spec}_X(\mathcal{O}_X/\text{cond}_X)$. Let $n : \bar{X} \rightarrow X$ be the normalization, and \bar{D} the pre-image of D in \bar{X} . Then there is an involution $\sigma : \bar{D}^n \rightarrow \bar{D}^n$ on the normalization of \bar{D} . We can write

$$n^*(K_X + \Delta) \sim_{\mathbb{Q}} K_{\bar{X}} + \bar{D} + \bar{\Delta},$$

where $\bar{\Delta}$ is the preimage of Δ . In fact, we only need to check this formula at all codimension 1 points, which is straightforward.

Definition 3.1. We call a demi-normal pair (X, Δ) is *semi-log-canonical* if $K_X + \Delta$ is \mathbb{Q} -Cartier and in the above notations, the pair $(\bar{X}, \bar{D} + \bar{\Delta})$ is log-canonical.

Definition 3.2. Let (X, Δ) be a demi-normal pair where $\Delta = \sum a_i \Delta_i$ is a sum of distinct prime divisors, none of which is contained in the singular locus $\text{Sing}(X)$ of X , and assume $0 \leq a_i \leq 1$ for every i .

We call a birational projective morphism $f : Y \rightarrow (X, \Delta)$ a *semi-log-canonical model* if f is isomorphic over open locus of X with complement's codimension greater than 1, and (Y, Δ_Y) is semi-log-canonical for $\Delta_Y = f_*^{-1}\Delta + E_f^{\text{lc}}$ where E_f^{lc} is the sum of all the exceptional prime divisors, and $K_Y + \Delta_Y$ is f -ample.

We note that from the definition, the induced map on the conductor schemes $D_Y \rightarrow D$ is an isomorphism outside some lower-dimensional subsets of D_Y and D , i.e., the codimension 1 points of the f -exceptional locus are all regular.

Lemma 3.1. *Given a demi-normal pair (X, Δ) , its semi-log-canonical model, if exists, is unique.*

Proof. Let Y be a semi-log-canonical model of (X, Δ) and $n_Y : \bar{Y} \rightarrow Y$ its normalization and $\bar{f} : \bar{Y} \rightarrow \bar{X}$ the induced morphism. We write

$$n_Y^*(K_Y + \Delta_Y) = K_{\bar{Y}} + \bar{D}_Y + \bar{\Delta}_Y.$$

Then $\bar{D}_Y + \Delta_Y = \bar{f}_*^{-1}(\bar{D} + \bar{\Delta}) + E_{\bar{f}}^{\text{lc}}$. Therefore, $\bar{f} : \bar{Y} \rightarrow (\bar{X}, \bar{D} + \bar{\Delta})$ is the log-canonical model, which is unique by (2.1). On a dense open subset of \bar{D}_Y , the involution $\sigma_Y : \bar{D}_Y \rightarrow \bar{D}_Y$ is the same as the restriction of $\sigma : \bar{D} \rightarrow \bar{D}$ to an isomorphic open subset, so σ_Y is uniquely determined, hence the quotient Y uniquely exists by [Kollár12, 5.3]. \square

On the other hand, with the results in [Kollár11] (also see [Kollár12, Section 4]), we can glue the log-canonical model of each component of the normalization $\bar{X} \rightarrow X$ to get the semi-log-canonical model of (X, Δ) .

Proof of (1.2). Let $f : \bar{Y} \rightarrow (\bar{X}, \bar{D} + \bar{\Delta})$ be the log-canonical model and write

$$f^*(K_{\bar{X}} + \bar{D} + \bar{\Delta}) = K_{\bar{Y}} + \bar{D}_Y + \bar{\Delta}_Y,$$

where \bar{D}_Y is the birational transform of \bar{D} on \bar{Y} . Then it follows from (2.1) that the normalization D_Y^n of \bar{D}_Y is the log-canonical model of (D^n, Δ_{D^n}) , where $K_{D^n} + \Delta_{D^n} = n^*(K_{\bar{X}} + \bar{D} + \bar{\Delta})|_{D^n}$ if we denote the normalization as $n : D^n \rightarrow D$.

Furthermore, because of the uniqueness of the log-canonical model by (2.1), this involution $\sigma : D^n \rightarrow D^n$ can be lifted to an involution on the log-canonical model as $\sigma_Y : D_Y^n \rightarrow D_Y^n$. Since $K_{\bar{Y}} + \bar{D}_Y + \bar{\Delta}_Y$ is ample over X , by [Kollár11, 26], $(\bar{Y}, \bar{D}_Y, \bar{\Delta}, \sigma_Y)$ has a quotient Y which is easy to see to be the semi-log-canonical model of (X, Δ) . \square

While the log-canonical models (1.1) are expected to exist even without the assumption that $K_X + \Delta$ is \mathbb{Q} -Cartier (see (2.1)), the next example constructed by Professor Kollár shows that in (1.2) the \mathbb{Q} -Cartier assumption on $K_X + \Delta$ is necessary. We are grateful to him for providing this example to us.

Example 3.1 (Kollár’s example on non-existence of semi-log-canonical models). We construct a demi-normal three-fold X with two irreducible components (X_i, D_i) such that X does not have an semi-log-canonical model. Take

$$(X_1, D_1) := (\mathbb{A}_{uvw}^3 / \frac{1}{3}(1, 1, 1), (w = 0) / \frac{1}{3}(1, 1) \cong \mathbb{A}_{uv}^2 / \frac{1}{3}(1, 1)).$$

(X_1, D_1) is lc (even plt), hence its log-canonical model is trivial, i.e., $\pi_1 : (Y_1, D_{Y_1}) \cong (X_1, D_1)$.

Note that $\mathbb{A}_{uv}^2 / \frac{1}{3}(1, 1)$ embeds in \mathbb{A}_{xyzt}^4 as the cone over the twisted cubic by $\sigma : (u, v) \mapsto (u^3, u^2v, uv^2, v^3)$; let $D_2 \subset \mathbb{A}_{xyzt}^4$ be its image. Then set

$$(X_2, D_2) := ((xt - yz = 0), D_2) \subset \mathbb{A}_{xyz}^4.$$

Use $\sigma : D_1 \cong D_2$ to glue (X_1, D_1) and (X_2, D_2) to obtain X .

To compute the log-canonical model over (X_2, D_2) , note that D_2 satisfies the equation $xz = y^2$ and $X_2 \cap (xz = y^2)$ is the union of D_2 and a residual plane $P := (x = y = 0)$.

Let $\pi_2 : Y_2 \rightarrow X_2$ be the blow up of the plane P and $C \subset Y_2$ the exceptional curve. Let D_{Y_2} (resp., \tilde{P}) denote the birational transforms D_2 (resp., P). Then $\pi_2^*(D_2 + P) = D_{Y_2} + \tilde{P}$ and $(C \cdot \tilde{P}) = \mathcal{O}(-1)|_{\mathbb{P}^1} = -1$. Thus $(C \cdot D_{Y_2}) = 1$ hence $K_{Y_2} + D_{Y_2}$ is π_2 -ample. By explicit computation, Y_2 and D_{Y_2} are both smooth, thus $\pi_2 : (Y_2, D_{Y_2}) \rightarrow (X_2, D_2)$ is the log-canonical model. Furthermore, $\pi_2 : D_{Y_2} \rightarrow D_2$ is the blow up of the origin, hence it is not an isomorphism. (We note that $K_{X_2} + D_2$ is not \mathbb{Q} -Cartier.)

Thus the isomorphism $\sigma : D_1 \cong D_2$ gives a birational map $\sigma' : D_{Y_1} \dashrightarrow D_{Y_2}$ that is not an isomorphism. Therefore (Y_1, D_{Y_1}) and (Y_2, D_{Y_2}) can not be glued together.

Acknowledgment

It is the authors’ great pleasure to thank Professors Christopher Hacon, Masayuki Kawakita, János Kollár and Shigefumi Mori for helpful e-mails and conversations. The joint work started when C.X. visited RIMS, he would like to thank the warm hospitality and inspiring environment there. Y.O. is partially supported by the Grant-in-Aid for Scientific Research (KAKENHI no. 21-3748) and the Grant-in-Aid for JSPS fellows. C.X. was partially supported by NSF research grant no. 0969495.

References

- [BCHM10] C. Birkar, P. Cascini, C. Hacon and J. Mckernan, *Existence of minimal models for varieties of log general type*, J. Amer. Math. Soc. **23**(2) (2010), 405–468.
- [Birkar11] C. Birkar, *Existence of log canonical flips and a special LMMP*, 2011, [arXiv:1104.4981](#).
- [Fujino10] O. Fujino, *Semi-stable minimal model program for varieties with trivial canonical divisor*, 2010, [arXiv:1010.2577](#).
- [Hacon11] C. Hacon, *On log canonical inversion of adjunction*, to appear in Proc. of Edin. Math. Soc., [arXiv:1202.0491](#).
- [HX11] C. Hacon and C. Xu, *Existence of log canonical closures*, 2011, [arXiv:1105.1169](#).
- [Kawakita07] M. Kawakita, *Inversion of adjunction on log canonicity*, Invent. Math. **167**(1) (2007), 129–133.
- [KM98] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Math., **134**, Cambridge University Press, Cambridge, 1998.
- [KK10] J. Kollár and S. Kovács, *Log canonical singularities are Du Bois*. J. Amer. Math. Soc. **23**(3) (2010), 791–813.



- [Kolláretal92] J. Kollár (ed), *Flips and abundance for algebraic threefolds*, *Astérisque* **211** (1992), 1–258.
- [Kollár12] J. Kollár, *Singularities of the minimal model program*. With the collaboration of S. Kovács. A book to appear.
- [Kollár11] J. Kollár, *Sources of log canonical centers*, 2011, [arXiv:1107.2863](https://arxiv.org/abs/1107.2863).
- [Odaka11] Y. Odaka, *The GIT stability of polarized varieties via discrepancy*, preprint, see <http://www.kurims.kyoto-u.ac.jp/~yodaka>.

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, JAPAN

E-mail address: yodaka@kurims.kyoto-u.ac.jp

URL: <http://www.kurims.kyoto-u.ac.jp/~yodaka>

BEIJING INTERNATIONAL CENTER OF MATHEMATICS RESEARCH, 5 YIHEYUAN ROAD, HAIDIAN DISTRICT, BEIJING 100871, CHINA

Current address: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, 155 SOUTH 1400 EAST, SALT LAKE CITY, UT 84112, USA

E-mail address: dr.chenyang.xu@gmail.com